



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Now assume

function $(b, c, d, e) = lb + mc + nd + pe)^2 + (qc + rd + se)^2 + (td + ne)^2 + (ve)^2$.

Expanding and equating to the corresponding coefficients of (4),

$l^2 = A_1$ (6); $lm = A_2$ (7); $ln = A_3$ (8); $lp = A_4$ (9); $m^2 + q^2 = A_5$ (10);
 $mn + qr = A_6$ (11); $mp + qs = A_7$ (12); $n^2 + r^2 + t^2 = A_8$ (13); $np + rs +$
 $nt = A_9$ (14); $p^2 + s^2 + u^2 + v^2 = A_{10}$. (15)

From (6) $l = A_1^{\frac{1}{2}}$; hence from (7), (8) and (9) we find m, n and p ; then from (10) we find q ; and from (11) and (12) we can then find r and s ; then from (13) and (14) we know t and u , and finally from (15) we find v . Thus equation (5) is always possible.

From $\Sigma(y^2) = (lb + mc + nd + pe)^2 + (qc + rd + se)^2 + (td + ue)^2 + (ve)^2 = 0$ we can assume $v^2e^2 = -(lb + mc + nd + pe)^2$ and $(td + ue)^2 = -(qc + rd + se)^2$, as the equation will then be satisfied. Therefore

$$ve = \sqrt{-1}(lb + mc + nd + pe), \quad (16)$$

$$td + ue = \sqrt{-1}(qc + rd + se). \quad (17)$$

From (16) and (17) we can express d and e by two equations of the *first* degree in b and c . Then, substituting in $\Sigma(y^3) = 0$, we have a homogeneous equation of the third degree in b and c . One of these may now be arbitrarily assumed and the other will then be found by the solution of an equation of the third degree.

If we make the above substitution in $\Sigma(y^4) = 0$ the final equation will be of the fourth degree. Therefore the general equation of the fifth degree can be reduced to either of the forms $x^5 + q_4x + q_5 = 0$ or $x^5 + q_3x^2 + q_5 = 0$.

In these two equations let $x = 1 \div z$. Substituting and reducing we get the two forms

$$z^5 + \frac{q_4}{q_5}z^4 + \frac{1}{q_5} = 0, \quad \text{and} \quad z^5 + \frac{q_3}{q_5}z^3 + \frac{1}{q_5} = 0.$$

This result was first obt'd by E. S. Bring and afterward by Mr. Jerrard.

Their method of making the substitutions may be seen in Todhunter's Theory of Equations or Serret's Cours D' Algebre' Superieure.

The above method is much easier than theirs, especially in finding the equation in y , which is there obtained by Tschirnhausen's substitution.

NOTE, BY MARCUS BAKER.—The Note on p. 53 of the last No. is an important one and I would very much like to see it made fuller.

Prob. 5, p. 66, Vol. I, has not been solved *in general*, the published solution applying only to a special case of the problem.

Prob. 49, p. 61, Vol. II, is erroneous and incomplete and needs further consideration.

Prob. 132, which you mention in your list, was completely solved by Prof. Hall at p. 55, Vol. IV. [This fact had escaped our recollection.—Ed.]